Assessment
- GCS
- V/S

↓

airway/ ventilation adequate — No → | PCP / ICP / ACP / CCP | see advanced airway management protocols |

Yes ↓

BP/ perfusion adequate — No → | PCP / ICP / ACP / CCP | see shock protocols |

Yes ↓

glucometer > 4mmol/L — No → | PCP / ICP / ACP / CCP | see hypoglycemia protocol |

Yes ↓

- abnormal speech - unilateral weakness - facial droop — Yes → **Determine 'Reperfusion Interval'*** — R.I.>3.5 Hours → **Transport to Nearest EHCF Hospital With CT**

No ↓

| PCP / ICP / ACP / CCP | **Transport** to nearest hospital |

(from Determine 'Reperfusion Interval'*) R.I.≤ 3.5 Hours ↓

| PCP / ICP / ACP / CCP | Transport to DASH (District Acute Stroke Hospital) |

↓

IMMEDIATELY CONTACT ED Physician ** At DASH (District Acute Stroke Hospital)

(from Transport to nearest hospital) ↓

Vomiting — Yes → | PCP / ICP / ACP / CCP | see nausea/ vomiting protocol |

No ↓

| ICP / ACP / CCP | IV saline lock |

* Reperfusion Interval (RI) = Duration of stroke symptoms plus the anticipated duration of transport to arrival at DASH (District Acute Stroke Hospital).